

## Predictors of Genital Pain in Young Women

Melissa A. Farmer  Cindy M. Meston

Received: 14 November 2005 / Revised: 14 September 2006 / Accepted: 28 October 2006  
Springer Science+Business Media, LLC 2007

**Abstract** Despite the high prevalence of genital pain in animals underlying the impairment of sexual function, healthy young adult women, limited research has addressed sexual behavior, and sexual satisfaction. genital pain during intercourse using contemporary models of multidimensional sexual function. The objectives of this study were threefold: (1) to identify differences in sexual functioning in women who experience genital pain compared to pain free women; (2) to identify predictors of sexual functioning in women with and without genital pain; and (3) to identify predictors of sexual satisfaction in women with and without genital pain. Sexually active female undergraduates ( $n = 651$ ) were administered the Female Sexual Function Index and the Derogatis Sexual Functioning Inventory. We evaluated the sexual factors that impact the sexual function of women with any pain (including high and low pain groups) versus women with no history of pain. Women with genital pain reported greater rates of sexual dysfunction as compared to pain free women; however, sexual functioning in the high versus low pain groups was distinguished primarily by vaginal lubrication. Women in the high pain group showed negative correlations between domains of sexual satisfaction and genital pain frequency and intensity that were not found in the low pain group. For pain-free women, underlying infection, allergies, muscle tension, hormone dysregulation played a strong role in sexual satisfaction, whereas non-intercourse sexual behavior was central to sexual satisfaction in women who reported pain. The evaluation of levels of genital pain may provide insight into the mechanism of localized or generalized genital contact, the most common type of dyspareunia is Vulvar Vestibulitis Syndrome (VVS, also known as vestibulodynia), or pain concentrated at the vaginal vestibule (Friedman, 1995).

**Keywords** Genital pain Dyspareunia Female sexual dysfunction Sexual satisfaction Sexual communication

---

M. A. Farmer  
Department of Psychology, McGill University, Montreal,  
Quebec, Canada

C. M. Meston & )  
Department of Psychology, University of Texas at Austin,  
1 University Station A8000, Austin, TX 78712, USA  
e-mail: meston@psy.utexas.edu

In addition to the physical pain, dyspareunia may result in a cascade of psychological, sexual, and interpersonal disturbances that impair a woman's quality of life and diminish satisfaction in primary intimate relationships (Bergeron, Binik, Khalife & Pagidas, 1997; Meana, Binik,

Khalife, & Cohen, 1997). The majority of our current knowledge about psychological factors in dyspareunia is based on premenopausal women with a mean age of 30 years (reviewed in Basson et al., 2004). Most evidence on dyspareunia in young adult women has been limited to samples of women with VVS (e.g., Bergeron et al., 2001; Danielsson, Sjoberg, & Wikman, 2000; Granot, 2005). Although VVS is thought to account for many cases of dyspareunia, it is unclear how these findings generalize to women with other types of dyspareunic pain. Subsequently, little is known about the experience of different levels of genital pain during early adulthood when dyspareunia is most prevalent (Laumann et al., 1999).

To date, the literature shows contradictory and inconclusive support for the contribution of psychological factors to the development and/or maintenance of dyspareunia. Aside from psychoanalysis, the literature on dyspareunia has often lacked theoretical direction and this may account for the disjointed psychological research. However, there is reason to expect a strong psychological component in the experience of genital pain. Psychological factors have been shown to impact the subjective experience of pain in a variety of pain populations (e.g., Vlaeyen & Linton, 2000). Because the experience of pain includes sensory and affective aspects, a woman's cognitive and emotional responses may exacerbate pain by increasing pelvic muscle tension and attention toward painful stimuli (Payne, Binik, Amsel, & Khalife, 2004).

An abundance of empirical studies and case reports have proposed a range of potential psychological risk factors for the development and maintenance of genital pain. Variables that have discriminated between healthy controls and women with dyspareunia include vulnerability factors, such as the personality trait neuroticism (e.g., Van Lankveld, Brewaeys, ter Kuile, & Weijenborg, 1995), mood disruptions (e.g., Gates & Galasko, 2001), underlying psychopathology, such as obsessive-compulsive traits and phobias (e.g., Meana et al., 1997; Van Lankveld et al., 1995), and individual difference factors like catastrophizing and hypervigilance (Payne et al., 2004; Pukall, Binik, Khalife, Amsel, & Abbott, 2002). However, many of these studies are limited by a lack of control groups and only a few psychological factors have been consistently addressed in the literature (e.g., depression, anxiety). Furthermore, many studies show conflicting support for these psychological contributions and it is unclear whether the psychological factors preceded pain or developed after the pain began (Basson et al., 2004).

Some women with dyspareunia report fear and disgust reaction to sexual intercourse (Kaneko, 2001). Importantly, affective responses, such as fear, can mediate the psychological response to pain, thereby resulting in increased attention to pain and thus greater perceived pain (Meagher, Arnau, & Rhudy, 2001). This fear may be due to a number of factors, including the anticipation of pain based on past painful sexual experiences, past sexual trauma, dysfunctional sexual schema (the mental framework with which one understands sexuality), or negative sexual attitudes. It is also possible that this fear predated the genital pain or has been maintained by persistent pain during sexual activity. When painful sexual experiences perpetuate fear and expectations of pain during sexual activity, a woman may choose to avoid sexual behavior altogether. Evidence supports many of these interpretations. Women with dyspareunia have self-reported more negative feelings surrounding sexual activity (Nunns & Mandal, 1997; Wouda et al., 1998), less sexual pleasure (Reissing, Binik, Khalife, Cohen, & Amsel, 2003), more feelings of depression about sexuality (Gates & Galasko, 2001), and negative sexual attitudes (Meana et al., 1997). Furthermore, one's sexual schema can be influenced by a number of sexual experiences, including age of sexual intercourse debut (Haavio-Mannila & Kontula, 1997; Sprecher, Barbee, & Schwartz, 1995), early non-intercourse sexual experience (Davis & Lay-Yee, 1999), number of sexual partners, exclusivity within a sexual relationship (Waite & Joyner, 2001), and importance of sex (Byers, 2005). Attitudes about gender-typical sexual behavior may also impact future sexual activity (Milhausen & Herold, 1999). Finally, it has been thought that women with dyspareunia are more likely to have experienced sexual abuse, although retrospective reports fail to support this assertion (see Dunne & Najman, 2005). In summary, little is known about what types of sexual experience may have contributed to the fear reported by women with dyspareunia.

In addition to psychological factors and fear of pain, the interpersonal impact of dyspareunia may be related to the quality of a woman's relationship with her partner and the accompanying sexual satisfaction. Existing evidence suggests that sexual satisfaction of women with dyspareunia is lower in comparison with control women (Danielsson et al., 2000; Wouda et al., 1998). Sexual satisfaction in healthy women is positively associated with young age, liberal attitudes toward sexuality, frequency of sexual intercourse (Renaud & Byers, 1997), relationship satisfaction (Lawrance & Byers, 1995; Sprecher, 2002), low discrepancy in sexual desire between partners (Davies, Katz, & Jackson, 1999), minimal cognitive distraction during sexual activity (Dove & Wiederman, 2000), self-disclosure, and masturbation (Byers & Demmons, 1999; Haavio-Mannila & Kontula, 1997; Sprecher & McKinney, 1993). Sexual satisfaction is strongly positive correlated with sexual communication, even when a woman's sexual pleasure is reduced (Bridges, Lease, & Ellis, 2004; Byers & Demmons, 1999). Women who are sexually dissatisfied may seek sexual expression through means independent of

intercourse with a primary partner, such as self-stimulation during masturbation and in the laboratory and or extra-pair sexual activity (Bridges et al. 2004; Davidson & Moore, 1994). Because women with dyspareunia tend to be younger (Laumann et al. 1999), engage in less sexual intercourse (Danielsson et al. 2000; Meana et al., 1997), report low sexual desire (Meana et al. 1997), and describe more fear associated with sexual activity (Kane et al. 2001), their understanding of sexual satisfaction may differ from women who do not experience pain during sexual intercourse. Due to the lack of evidence about how genital pain affects the experience of sexual satisfaction, it would be useful to evaluate how the factors that predict levels of sexual satisfaction in women with dyspareunia differ from those found in pain-free women.

Another consequence of the psychological impact of dyspareunia is the potential for coexisting sexual dysfunction. Importantly, inhibited desire, arousal, and lubrication are thought to disrupt the psychological and physiological progression of the sexual response cycle and may result in greater genital pain (Basso 2001). This disruption may occur through the anticipation of pain, which may lead to cognitive responses that suppress sexual response. For instance, the expectation of pain may result in self-monitoring during sexual activity, and self-monitoring is known to inhibit the sexual response (Dove & Wiedermayr 2000; Van den Hout & Barlow, 2000). Indeed, women with dyspareunia may experience lower sexual desire (Meana et al., 1997; Reissing et al. 2003), arousal (e.g., Reissing et al., 2003; Wouda et al., 1998), and lubrication (Nunns & Mandal, 1997; Van Lankveld, Weijenborg, ter Kuile, 1996). In a sample of women with vulvar pain, the impaired sexual arousal disorder, with the addition of increased pain and lower lubrication (Masheb, Lozano-Blanco, Kohorn, Minkin, & Kerns, 2004). An inverse relationship between genital pain and lubrication has since been reported (Wiegand, Meston, & Roser 2005). However, the extent of sexual dysfunction in this population is a point of contention. Two independent studies measuring genital arousal in response to erotic video stimuli found that women with dyspareunia did not differ from controls in levels of vaginal blood flow or in subjective reports of genital arousal (Brauer, Laan, & ter Kuile, 2006; Wouda et al., 1998). Similarly, Van Lankveld et al. (1996) reported no significant differences in self-reported subjective arousal, lubrication, and genital pain during masturbation in a sample of women with VVS.

In contrast to masturbation, self-reported sexual problems and distress during sexual intercourse were significantly greater in these women. The ability of some women with dyspareunia to achieve a normal sexual response in some contexts suggests that sexual dysfunction is not a necessary or sufficient requirement for genital pain. It is possible that women with dyspareunia are capable of a normal sexual

response during masturbation and in the laboratory setting, we predicted that women with high levels of pain would report equal levels of non-intercourse sexual behavior (including masturbation, petting, and oral sex) and lower levels of intercourse as compared to women without pain.

Method  
Participants  
Female undergraduates ( $n = 871$ ) from a large public university participated in exchange for course credit between 2001 and 2004. Cohorts included participants from spring and fall semesters and varied in size (2001  $n = 248$ ; 2002,  $n = 165$ ; 2003  $n = 236$ ; 2004  $n = 222$ ). Primary inclusion criteria was sexual activity within one month of participation, and the revised sample ( $n = 651$ ) was used for final analyses.

All participants were between 18 and 25 years old ( $M = 18.72$ ). A high percentage of this sexually active sample had previously experienced sexual intercourse (93%), and a slightly smaller percentage (70%) reported involvement in a long-term relationship in the 12 months before participation. The sample consisted of predominately Caucasian participants, with 64% Caucasian (486), 17%

1. Identify differences in sexual function, satisfaction, behavior, and attitudes in women with no, low, and high pain during sexual intercourse.  
2. Identify the factors that statistically predict the sexual function of sexually active young adult women who have and have not reported genital pain.  
3. Identify the statistical predictors of sexual satisfaction in sexually active young women who have and have not reported genital pain.

1. Identify differences in sexual function, satisfaction, behavior, and attitudes in women with no, low, and high pain during sexual intercourse.
2. Identify the factors that statistically predict the sexual function of sexually active young adult women who have and have not reported genital pain.
3. Identify the statistical predictors of sexual satisfaction in sexually active young women who have and have not reported genital pain.

In line with past research, we hypothesized that women with any genital pain were more likely to report poor sexual function, more conservative sexual attitudes, and less sexual satisfaction when compared with pain-free women. Due to past findings that women with dyspareunia have exhibited normal sexual arousal during masturbation and in the laboratory setting, we predicted that women with high levels of pain would report equal levels of non-intercourse sexual behavior (including masturbation, petting, and oral sex) and lower levels of intercourse as compared to women without pain.

Method  
Participants  
Female undergraduates ( $n = 871$ ) from a large public university participated in exchange for course credit between 2001 and 2004. Cohorts included participants from spring and fall semesters and varied in size (2001  $n = 248$ ; 2002,  $n = 165$ ; 2003  $n = 236$ ; 2004  $n = 222$ ). Primary inclusion criteria was sexual activity within one month of participation, and the revised sample ( $n = 651$ ) was used for final analyses.

All participants were between 18 and 25 years old ( $M = 18.72$ ). A high percentage of this sexually active sample had previously experienced sexual intercourse (93%), and a slightly smaller percentage (70%) reported involvement in a long-term relationship in the 12 months before participation. The sample consisted of predominately Caucasian participants, with 64% Caucasian (486), 17%

123

Hispanic ( $n = 125$ ), 13% Asian American ( $n = 100$ ), 4% African American ( $n = 27$ ), 3% other ethnicities (21) participants. Represented religious groups include Atheist (13%), Agnostic (13%), Buddhist (2%), Hindu (4%), Jewish (5%), Protestant (16%), Catholic (26%), Fundamentalist Christian (33%), and Muslim (1%) participants.

Pain groups were categorized using an index score that reflected frequency and intensity of genital pain during and after intercourse (see Sexual Functioning for a detailed description of this index). Sixty three percent of sexually active women reported any history of genital pain during intercourse. Of sexually active women, 40% of women ( $n = 217$ ) reported genital pain occasionally or sometimes, women who reported never having experienced pain during intercourse were classified as having no history of pain during sexual intercourse. The pain domain time. No significant demographic differences were found between women who reported no history of genital pain and women who reported a history of genital pain.

## Measures

### Sexual functioning

Sexual functioning was assessed using the Female Sexual Functioning Index (FSFI; Rosen et al., 2000). The FSFI is a brief, 19-item scale, which measures extent of difficulty in six domains of sexual function: desire (two items), arousal (four items), lubrication (four items), orgasm (two items), sexual satisfaction (three items), and pain (three items). Items used a response format based on the response range of one (almost never or never) to five (almost always or always). Higher FSFI scores are associated with more healthy sexual functioning. Psychometrically, the FSFI has robust reliability and validity across populations of sexually healthy and dysfunctional women, including women with vulvodynia and women with symptoms of dyspareunia (Masheb et al., 2004; Meston, 2003; Wiegel et al., 2005). The scale has shown appropriate internal reliability ( $\alpha = .89$ -.97) and test-retest reliabilities (Cronbach's alpha = .79-.88). In the current sample, reliability coefficients were within an acceptable range for the domains of desire (Cronbach's alpha = .86), arousal (Cronbach's alpha = .94), lubrication (Cronbach's alpha = .95), orgasm (Cronbach's alpha = .92), and satisfaction (Cronbach's alpha = .84). A summed FSFI total score, which did not include the pain domain, was used to measure sexual function. In addition to assessment of sexual function, the FSFI was used to classify participants into currently sexually active and inactive groups.

The FSFI pain domain score was used to create no, low, and high groups for primary analyses. The pain domain included three questions: "How often do you experience pain during penetration" included responses from one (almost always or always) to four (almost never or never),

"How often do you experience pain after penetration?" included a response format of one (most times) to five (almost never or never); and "Rate your level of pain during or following penetration" included a response format of one (very high) to five (very low or none). Lower scores indicated greater pain during sexual intercourse.

Scores were used to classify women into three groups: no pain, low pain, and high pain. Women who reported pain "always" or "almost always" were coded as the high pain group (FSFI pain subscore  $\leq 3.20$ ). Women who "occasionally" or "sometimes" experienced pain were coded as the low pain group (pain subscores between 3.60 and 5.20). Finally, women who reported "never" having experienced pain during intercourse were classified as having no history of pain (pain subscores over 5.20). The pain domain showed excellent reliability coefficients in the current sample (Cronbach's alpha = .97).

### Sexual satisfaction

Sexual satisfaction was evaluated with the Sexual Satisfaction Scale for Women (SSS-W; Meston & Trapnell, 2005), which consists of 30 Likert-type items that describe five domains of sexual satisfaction: Personal Concern (e.g., "I'm so distressed about my sexual difficulties that it affects the way I feel about myself"), Interpersonal Concern (e.g., "I feel like I've disappointed my partner by having sexual difficulties"), Compatibility (e.g., "I sometimes think my partner and I are mismatched in needs and desires concerning sexual intimacy"), Communication (e.g., "I usually feel completely comfortable discussing sex whenever my partner wants to"), and Contentment (e.g., "I feel content with the way my present sex life is"). Response options ranged from one (strongly disagree) to five (strongly agree), and questions with negative phrasing were reverse scored so that higher numbers reflect greater sexual satisfaction. For the current sample, reliability coefficients were adequate (Cronbach's alpha = .89).

### Unrestricted sexual behavior

Items from the Sociosexual Orientation Inventory (SOI; Simpson & Gangestad, 1991) assessed sociosexual behavior and attitudes. The SOI was designed to measure individual differences in the endorsement of restricted sexual behaviors and attitudes (i.e., restricting sexuality to intimate and committed contexts) or unrestricted sexual behavior and attitudes (i.e., sexuality is not restricted to intimacy and commitment and may be more casual in nature). Unrestricted sexual behavior includes the number of sexual intercourse partners within the past year, number of one-time instances of sexual intercourse ("One night

stands (Ö), and anticipated number of sexual partners in Results following 5 years. In addition to SOI questions, the authors

included two related unrestricted sexuality items. Participants endorsed the number of individuals with whom they had engaged in foreplay within the past year and the lifetime number of individuals with whom they had engaged in sexual intercourse. The sample reliability for the Pve items was acceptable (Cronbach's alpha = .83).

### Sexual attitudes and behavior

The Sexual Attitude and Sexual Experience subscales of the Derogatis Sexual Functioning Inventory (DSFI; Derogatis & Melisaratos, 1979) were used to assess a variety of sexual functioning domains. The DSFI Sexual Attitude Scale consists of statements reflecting a continuum of conservative to liberal sexual attitudes. Items were coded so that higher numbers reflected more conservative sexual attitudes.

The DSFI Experience Scale was used to assess the following sexual experience domains: petting (six items), oral sex (Pve items), intercourse (four items), and masturbation (one item). Using a yes/no response format, participants indicated if they had ever engaged in the specified sexual activity. Coding procedures assigned yes (1) or no (0) values to each domain, and group averages were obtained. The Experience Scale has shown high internal consistency (.97) and test-retest reliability (.92).

### Procedure

Female researchers administered questionnaires to groups of 5–10 participants. Testing took place in unoccupied computer laboratories and classrooms, and participants were separated by a minimum of 5 feet to maximize privacy. All participants were aware of the sensitive nature of the questionnaire material before testing. The study procedure was explained and participants were encouraged to ask questions or raise concerns before giving informed consent. It was emphasized that if participants experienced discomfort or distress during testing, then they were encouraged to pause or stop participation. The researcher informed the testing group that participation was completely voluntary, that anonymity and confidentiality of data would be maintained, and that no future relationship with the institution would be adversely impacted by refusal to participate. No participants chose to stop participation. Data were locked in a filing cabinet accessible only to the primary investigators. Electronic data files were password protected.

Differences between the pain groups and healthy controls are reported in columns 2–4 of Tables 3, and 4. In order to test our hypotheses, analyses were conducted on two subgroups: women who reported no pain and women who reported pain. When significant main effects were found, the Games-Howell post-hoc test for unequal group sizes and unequal variances was performed on the no, low, and high groups. To minimize the risk of Type I error in multiple comparisons, Bonferroni corrections indicate statistically reliable differences and are noted at the bottom of each table.

Main effects of no, low, and high pain groups were found for sexual function variables ( $p < .05/5$ ). The pain groups reported more difficulties with drive,  $F(2, 545) = 4.52, p < .01$ , arousal,  $F(2, 545) = 8.14, p < .01$ , lubrication,  $F(2, 545) = 184.60, p < .01$ , orgasm,  $F(2, 545) = 13.77, p < .01$ , and satisfaction,  $F(2, 545) = 5.60, p < .01$ . As indicated by the subscripts in Table 1, the Games-Howell post-hoc analyses indicated that the no pain group reported significantly better sexual function than the low and high pain groups. Additionally, the low pain group reported higher lubrication and overall sexual function scores as compared to the high pain group. No other significant sexual function differences were found between the low and high pain groups. Results indicated that the pain groups reported higher rates of sexual dysfunction within the previous month than the no pain group, and the high pain group reported significantly greater overall dysfunction and less lubrication than the low pain group.

No significant main effect was found for the SSS-W sexual satisfaction summed score (Table 1). The significant intercorrelations between sexual satisfaction variables were expected because they represent closely related constructs. However, the correlations between sexual satisfaction variables (contentment, communication, compatibility, interpersonal concern, and personal concern) and individual pain items were different for the low versus high pain groups (Table 2). In the low pain group, the only significant correlations found were between frequency of pain after penetration and communication ( $r = .15, p = .03$ ) and frequency of pain during/after penetration and frequency of pain during penetration ( $r = .18, p = .01$ ). In contrast, in the high pain group the frequency of pain during penetration was negatively correlated with communication ( $r = -.29, p < .01$ ), compatibility ( $r = -.22, p = .02$ ), and interpersonal concern ( $r = -.24, p = .01$ ). Frequency of pain after penetration was positively correlated with frequency of pain during/after penetration was negatively associated with contentment ( $r = -.20, p = .02$ ), compatibility ( $r = -.24,$

Table 1 Differences in sexual function and satisfaction between women with no pain, low pain, and high genital pain

Items	Women reporting pain								F value
	No pain		Any pain <sup>f</sup>		Low pain		High pain		
	M	SD	M	SD	M	SD	M	SD	
FSFI subscales									
Drive	4.48 <sup>g</sup>	1.02	4.21	1.02	4.24	1.00	4.17 <sup>h</sup>	1.06	4.52*
Arousal	5.13 <sup>g</sup>	.98	4.78	.99	4.75	.96	4.82 <sup>h</sup>	1.10	8.14*
Lubrication	5.47 <sup>h</sup>	.74	4.55	1.33	5.17 <sup>h</sup>	.87	3.49 <sup>h</sup>	1.33	184.60*
Orgasm	4.40 <sup>h</sup>	1.55	3.80	1.36	3.92	1.43	3.58 <sup>h</sup>	1.21	13.77*
Satisfaction	4.95 <sup>n</sup>	1.25	4.59	1.26	4.64	1.18	4.52 <sup>h</sup>	1.40	5.60*
FSFI total score <sup>e</sup>	24.44 <sup>p</sup>	4.08	21.94	3.88	22.72	3.77	20.59 <sup>q</sup>	3.70	38.74*
Sexual satisfaction total score	109.36	21.99	106.29	20.93	106.18	20.13	106.45	22.09	0.93
	n = 140 <sup>D</sup> 205		n = 200 <sup>D</sup> 293		n = 100 <sup>D</sup> 137		n = 65 <sup>D</sup> 79		

\*p < .01 based on a one-way ANOVA comparison between no pain, low pain, and high pain groups

<sup>a</sup> FSFI total excludes pain subscale and values range from 5 (poor sexual function) to 30 (high sexual function)

<sup>b</sup> SSS-W contains contentment, communication, compatibility, interpersonal concern, and personal concern subscales rated on a scale of 1 (strongly agree) to 5 (strongly disagree)

<sup>c</sup> The any pain group is further divided into the low and high pain groups

<sup>d</sup><sup>e</sup><sup>q</sup> Means with the same subscript differ significantly at .05

Table 2 Correlations between sexual satisfaction variables, frequency of genital pain, and level of genital pain

	1	2	3	4	5	6	7	8
1. Contentment	∅							
2. Communication	.53**	∅						
3. Compatibility	.57**	.53**	∅					
4. Interpersonal concern	.67**	.57**	.53**	∅				
5. Personal concern	.72**	.59**	.57**	.63**	∅			
6. Frequency of pain during penetration	.47**	.48**	.47**	.63**	.78**	∅		
7. Frequency of pain after penetration	.65**	.45**	.70**	.63**	.78**	.05	∅	
8. Level of pain during or following penetration	.48**	.40**	.52**	.63**	.78**	.05	.09	∅
	.70**	.42**	.68**	.78**	.78**	.05	.20*	.13
	.03	.00	.07	.05	.05	.05	.09	.13
	∅.15	∅.28**	∅.22**	∅.14	∅.24*	∅.05	∅.09	∅.13
	.04	.15*	∅.06	.02	∅.05	∅.05	∅.09	∅.13
	∅.04	∅.14	∅.06	∅.14	∅.08	∅.08	.20*	.13
	∅.01	.04	∅.08	.07	∅.11	∅.11	.18**	.13
	∅.20*	∅.17	∅.24**	∅.21*	∅.31**	∅.31**	.50**	.35**

\*p < .05, \*\*p < .01, the low pain group coefficients are listed above the high pain group coefficients

SSS-W items have response format of 1 (strongly disagree) to 5 (strongly agree). Questions are scored so that higher numbers reflect greater sexual satisfaction

p = .01), interpersonal concern (r = ∅.21, p = .03), and personal concern (r = ∅.31, p = .001) and positively correlated with frequency of pain during (r = .50, p < .01) and after (r = .35, p < .01) penetration. These findings suggest that pain variables in the low pain group were minimally related to sexual satisfaction, whereas the pain variables in the high pain group were often related to sexual satisfaction. No significant main effects were found for sexual experience variables (masturbation, petting, oral sex, and intercourse) or for unrestricted sexuality variables (p < .05/9). The majority of women from the pain and no pain groups reported experience with multiple types of sexual behavior (see Table 3 for percentages and behavior means). Sexual attitudes were evaluated in Table 4 (p < .05/14). Women with a history of pain endorsed significantly more

Table 3 Differences in sexual experience and unrestricted sexual behavior between women with no, low, and high genital pain

Items	Women reporting pain								F value
	No pain % Yes		Any pain <sup>c</sup> % Yes		Low pain % Yes		High pain % Yes		
	M	SD	M	SD	M	SD	M	SD	
Petting experience <sup>a</sup>	98		98		97		100		0.93
Oral sex experience <sup>a</sup>	94		96		96		97		0.46
Intercourse experience <sup>a</sup>	82		87		86		91		2.29
Masturbation experience <sup>a</sup>	75		73		74		71		0.24
No. foreplay partners in past year	3.66	3.62	4.07	4.06	3.93	4.00	4.31	4.16	1.10
No. intercourse/oral sex partners in past year	1.94	1.87	2.27	2.25	2.20	2.36	2.38	2.33	1.70
No. intercourse/oral sex partners just once	1.37	2.28	1.51	2.01	1.37	1.86	1.75	2.23	1.48
No. intercourse/oral sex partners lifetime	4.33	4.47	4.67	5.16	4.56	4.89	4.86	5.61	0.44
No. anticipated intercourse/oral sex partners	3.26	3.75	3.79	4.42	3.51	3.87	4.27	5.19	2.30
	n = 107		n = 212		n = 93		n = 64		

<sup>a</sup> Sexual experience items based on a yes/no response format

<sup>b</sup> Values represent the number of sexual partners endorsed by participants

<sup>c</sup> The any pain group was further divided into the low and high pain groups

conservative attitudes about whether it is unnatural for a woman to initiate sexual relations. In contrast, the group reported more liberal attitudes about whether anything is wrong with mate swapping.

#### Hierarchical multiple regressions

Two sets of hierarchical multiple regression analyses were conducted to statistically predict sexual function and sexual satisfaction in sexually active women who reported no history of genital pain and any history of genital pain so that individual pain variables could be evaluated. In the first set of regressions, women without a history of pain, the summed FSFI score (excluding the pain domain) was entered as the dependent variable, and the summed SSS-W sexual satisfaction score was entered as the dependent variable in the second set of regressions. In all regressions, a hypothesized model of variables linked to dyspareunia was entered as independent variables in each regression (including sexual function, sexual behavior during intercourse, women with healthy sexual history, and sexual satisfaction). The order in which variables were entered was based on the assumption that developmental, behavioral, sexual functioning, and psychological variables uniquely contribute to variance in penetration, frequency of pain after current sexual function. Therefore, Step 1 of the hierarchical regressions included age and sexual function, Step 2 included sexual behavior and sexual experience, and Step 3 included sexual satisfaction and psychological variables. For each sample, correlations were calculated between the dependent variable and the hypothesized model of predictors, and only variables that were significantly correlated with the dependent variable were entered into subsequent regressions. For each regression, multicollinearity diagnostic

Table 4 Differences in sexual attitudes between women with and without genital pain

Items	Women reporting pain								F value
	No pain		Any pain <sup>a</sup>		Low pain		High pain		
	M	SD	M	SD	M	SD	M	SD	
Sex without love is okay (R).	3.47	1.45	3.44	1.45	3.47	1.42	3.38	1.52	.08
Premarital intercourse is beneficial for later marital adjustment (R).	2.71	1.26	2.64	1.20	2.65	1.15	2.62	1.30	.52
Homosexuality is perverse and unhealthy.	2.34	1.30	2.43	1.43	2.43	1.38	2.43	1.51	.53
Oral sex can be as pleasurable as intercourse (R).	2.11	1.21	2.15	1.19	2.19	1.24	2.09	1.11	.15
It is unnatural for the woman to be the initiator of sexual relations.	1.60	0.90	1.89	1.11	1.91	1.09	1.86	1.15	6.32*
Masturbation is normal and healthy (R).	1.93	1.17	2.00	1.15	2.08	1.19	1.87	1.08	.49
Extramarital sex inevitably leads to marital problems.	4.17	1.22	4.05	1.26	3.94	1.30	4.25	1.17	1.13
Viewing erotic films is enjoyable and stimulating behavior (R).	2.39	1.18	2.57	1.28	2.52	1.25	2.63	1.32	2.57
There is nothing wrong with mating swapping (R).	4.36	1.07	4.13	1.17	4.12	1.12	4.14	1.25	5.51*
Males lose respect for females who have premarital sex.	2.42	1.30	2.33	1.20	2.30	1.21	2.36	1.19	.82
Group sex is bizarre and disgusting.	3.40	1.44	3.49	1.41	3.41	1.40	3.64	1.42	.57
Extramarital affairs can make people better partners (R).	4.50	0.91	4.36	1.00	4.37	0.94	4.35	1.09	2.75
I can imagine feeling comfortable with casual sex (R).	4.01	1.29	3.75	1.34	3.77	1.33	3.71	1.36	4.97
Homosexuality is not good or bad (R).	2.21	1.36	2.20	1.39	2.18	1.35	2.24	1.45	.00
	n = 203		n = 205		n = 342		n = 344		
	n = 216		n = 217		n = 126		n = 127		

\*p < .05, \*\*p < .01. Based on a response format of 1 (completely disagree) to 5 (completely agree) such that higher numbers indicate more conservative sexual attitudes. Reversed (R) items reflected a response format of 5 (completely agree) to 1 (completely disagree)

<sup>a</sup> The any pain group is further divided into the low and high pain groups

<sup>b</sup><sup>c</sup> Means with the same subscript differ significantly at .01

the second step. The hypothesized model ( $F(3, 143) = 4.65$ ,  $\eta^2 = .23$ ,  $p < .01$ ). These significant correlates of  $p < .01$ , indicated that 19% of the variance in sexual sexual satisfaction were entered into a regression, with functioning was explained by levels of pain frequency sexual function variables entered into the first step, sexual during penetration, pain frequency following penetration, behavior variables entered into the second step, and mas- level of pain during or after penetration, likelihood of turbation attitudes entered into the third step. The resulting cheating if totally committed, likelihood of cheating if not model,  $F(3, 132) = 9.09$ ,  $p < .01$ , accounted for 33% of the totally committed, overall sexual satisfaction, and tradi-variance in levels of sexual satisfaction in women who tional gender role attitudes. Unique contributions were reported no history of genital pain. Unique contributions to provided by overall sexual satisfaction  $\beta = .27$ ,  $t = 3.39$ , the model were made by the FSFI arousal domain  $\beta = .25$ ,  $p < .01$ ) and likelihood of cheating if totally committed  $t = 2.10$ ,  $p < .05$ ) and the intercourse subscore  $\beta = -.34$ ,  $(\beta = -.34, t = -2.61, p < .01)$ . The model indicated that in  $t = 4.12, p < .01$ ). The model suggested that in women who women with any history of genital pain during intercourse, report no history of genital pain during intercourse, women women with high levels of sexual functioning are more who have high sexual satisfaction are more likely to report likely to report greater sexual satisfaction and were less high levels of sexual arousal and are more likely to have likely to report they would cheat if totally committed to a experience with sexual intercourse (see Table partner. All beta values are summarized in Table

In women with any history of pain, an additional A second set of regressions was then computed to predict regression was calculated to predict sexual satisfaction sexual satisfaction in sexually active women with and (Table 7). Variables significantly correlated with sexual without a history of pain. In women with no history of pain, satisfaction in sexually active women with any history of sexual satisfaction was significantly correlated with the pain included the FSFI arousal domain  $(.21, p < .01)$ , FSFI arousal domain  $(.35, p < .01)$ , the FSFI lubrica- the FSFI orgasm domain  $(.26, p < .01)$ , the petting tion domain  $(.21, p < .05)$ , the FSFI orgasm domain subscore  $(.19, p < .01)$ , the oral sex subscore  $\beta = .27$ ,  $(r = .28, p < .01)$ , oral sex experience subscore  $(.21, p < .01)$ , number of sexual foreplay partners in the previous  $p < .01$ , intercourse experience subscore  $(.40, p < .01)$ , year  $(t = -1.16, p < .05)$ , number of sexual intercourse number of sexual foreplay partners in the previous year partners in the previous year  $\beta = -1.15, p < .05$ , anticipated  $(r = -1.29, p < .01)$ , anticipated number of sexual partners in number of sexual partners in the next 5 years  $\beta = -1.25$ , the next 5 years  $(\beta = -1.25, p < .01)$ , and masturbation atti-  $p < .01$ ), lifetime number of sexual partners  $\beta = -1.16$ ,

Table 5 Regressions to predict sexual function in women who report genital pain

Predictor Variables	$\hat{R}$	F	$\beta$	t	p
Model for total sample	.19	4.65**			
Frequency of pain during penetration				.14	1.16 ns
Frequency of pain after penetration				Đ.19	Đ1.48 ns
Pain level during/after sexual intercourse				.13	.91 ns
Sexual satisfaction				.27	3.39 .001
Likelihood of cheating if not totally committed to partner				.05	.41 ns
Likelihood of cheating if totally committed to partner				Đ.34	Đ2.61 .01
Gender role attitudes				Đ.02	Đ.19 ns

\*\* p < .01, sexual function measured with FSFI total score without pain domain

Table 6 Regressions to predict sexual satisfaction in women who report no genital pain

Predictor Variables	Adj. $\hat{R}$	F	$\beta$	t	p
Model for total sample	.33	9.09**			
Arousal				.25	2.10 .037
Lubrication				Đ.04	Đ.38 ns
Orgasm				.07	.73 ns
Oral sex experience				.07	.86 ns
Intercourse experience				.34	4.12 .001
No. foreplay partners				Đ.17	Đ1.88 ns
No. anticipated partners in next 5 years				Đ.15	Đ1.61 ns

\*\* p < .01, sexual satisfaction measured with the SSSDW

p < .05), number of one-time sexual partners = (Đ.21, likely to have experience with oral sex, and expected a p < .01), and masturbation attitudes = (.16, p < .05). lower number of sexual partners within the next 5 years. Sexual function variables were entered into the first step of the regression, sexual behavior variables were entered into the second step, and the third step included masturbation attitudes. The model F(2, 168) = 6.04 p < .01, explained 25% of the variance in sexual satisfaction, and unique contributions to the model were made by the FSFI orgasm behavioral, and attitudinal differences among women with domain ( $\beta = .26$ , t = 3.08, p < .01), oral sex experience no, low, and high genital pain. Additionally, the study subscale ( $\beta = .27$ , t = 2.46, p < .05), and anticipated number of sexual partners over the next 5 years = (Đ.17, sexual satisfaction in women with and without pain. As t = Đ2.00, p < .05). Of sexually active women with a history of genital pain, women with high sexual satisfaction significantly greater impairments in sexual function compared to women with no pain. The low and high pain groups

Table 7 Regressions to predict sexual satisfaction in women who report genital pain

Predictor Variables	Adj. $\hat{R}$	F	$\beta$	t	p
Model for total sample	.25	6.04**			
Arousal				.04	.50 ns
Orgasm				.26	3.08 .002
Petting experience				.07	.61 ns
Oral sex experience				.27	2.46 .015
No. foreplay partners				Đ.06	Đ.63 ns
No. intercourse partners				Đ.01	Đ.06 ns
No. anticipated intercourse partners				Đ.17	Đ2.00 .047
No. lifetime sexual partners				Đ.05	Đ.36 ns
No. one-time sexual partners				Đ.10	Đ.88 ns

\*\* p < .01, sexual satisfaction measured with the SSS-W

reported equally impaired desire, arousal, orgasm, and interpersonal concern and pain during sexual activity. Interestingly, women with low pain reported significantly more lubrication during sexual activity and reported higher global sexual functioning than women with high pain. These findings were consistent with past work showing a negative association between genital pain and lubrication, although it is unclear whether lubrication increases pain, pain reduces lubrication, or whether a third variable is driving the relationship (Wiegel et al., 2005).

It is interesting that the low pain group has retained some ability to maintain physiological sexual arousal despite reporting otherwise equivalent deficits in sexual function compared to the high pain group. Three interpretations may explain this phenomenon. One interpretation is that the presence of genital pain is correlated with the general impairment in sexual function. However, if the mere presence of pain during sexual activity is correlated with sexual dysfunction, then there would be little reason to believe a woman who experiences pain occasionally versus always would differ in function. An increase in the frequency and/or intensity of pain may then be associated with increased dysfunction appears to be specific to lubrication. A second interpretation is that lubrication is the functional deficit that distinguishes between high and low genital pain. Lack of lubrication can be both a cause and a consequence of painful intercourse (Binik, Wiegel et al., 2005). In the future, it would be useful to assess differences in lubrication between vaginal penetration and solitary self-stimulation by means other than self-report that baseline lubrication can be compared to levels achieved during intercourse (Van Lankveld et al., 1996). The third interpretation is that the low and high pain groups differ in etiology and their different patterns of sexual problems reflect this difference. If this difference in sexual response is indicative of a difference in the quality or type of pain experienced by the two groups, it is feasible that this sample the mechanisms underlying genital pain are different for the low and high pain groups.

Meana, Binik, Khalife and Cohen (1999) found that perceived differences in etiology were related to self-reported levels of pain. Women who believed their genital pain to be of psychosocial origin reported greater sensory pain and more intense experiences of pain as compared to women who attributed their dyspareunia to physical causes. The women who made psychosocial attributions were also more likely to report sexual problems, including more sexual aversion and less self-reported sexual arousal during intercourse and sexual satisfaction with a partner. It is possible that the high pain group may preferentially represent women who make psychosocial attributions. Such an explanation could account for the correlation between personal attribution and sexual satisfaction. In contrast, the pain experience of a new partner (Greiling & Busby, 2000). It would be useful to understand what motivations underlie the sexual activity perceived by women with genital pain (Hill & Prestor, 1996). Regardless of sexual motivation, it is clear that the experience of pain does not appear to prevent young adult women from engaging in sexual activity. In contrast to past research (e.g., Gates & Galanter, 1981), sexual satisfaction did not differ between women with pain and healthy women. Women with high pain reported a significant negative association between pain frequency and sexual satisfaction with a partner. It seems that for women with high levels of genital pain, the significance of intercourse may be difficult to escape when it comes to sexual satisfaction. In contrast, the pain experience

rience was not a salient component of sexual satisfaction in women with low levels of pain. This finding is reminiscent of Meana et al. (1998) finding that levels of marital adjustment predicted pain ratings in women with dyspareunia, such that women who endorsed high levels of marital adjustment were more likely to report poorer marital adjustment in the Locke-Wallis Marital Adjustment Scale. Notably, this measure of marital adjustment included sexual satisfaction.

For women who reported genital pain, sexual satisfaction was predicted by improved quality of orgasm, greater oral sex experience, and fewer projected future sexual partners. These findings are intriguing for two reasons. First, women with low and high pain reported impairments in multiple domains of sexual function, yet the majority of these sexual problems did not play a significant role in their perceptions of sexual satisfaction. Indeed, the finding that high sexual function was predicted by greater sexual satisfaction in women with pain suggests that a woman's comfort with sex and the interpersonal effects of genital pain. Finally, the interpersonal dynamics of sexual activity rather than the mere presence of pain is closely related to her ability to subjectively and physiologically respond to sexual stimulation. Second, rates of sexual intercourse were not related to sexual satisfaction in women who had experienced genital pain, whereas intercourse was a predictor of sexual satisfaction for women without pain. Instead, the sexual satisfaction of women with pain was predicted by oral sex experience, a non-intercourse behavior. These findings suggest that optimal levels of sexual satisfaction are independent of intercourse behavior in women with genital pain. Ironically, the frequency of sexual intercourse has been the primary measure of sexual disturbance in women with dyspareunia and has been the target of most interventions. It is possible that intercourse frequency has much less to do with the quality of a woman's sexual life than the practice of other sexual activity that is usually relegated to the realm of foreplay.

The prevalence of frequent genital pain during intercourse in the current sample was surprisingly high, particularly when women were not recruited based on sexual functioning variables. Additional information is needed in order to understand what factors may contribute to this high prevalence, including use of medications that alter sexual function (e.g., antidepressants), use of oral contraceptives, and contextual factors (type of stimulation received, activities that evoke pain, whether the pain is partner-specific). Partner characteristics, such as limited knowledge/use of sexual technique or premature ejaculation, may also influence a woman's experience of genital pain. The impairment in sexual function reported by women with pain suggests that these women experience difficulties with multiple aspects of the sexual response cycle. The reported pain could then be secondary to a variety of desire or arousal disorders.

Limitations that should be considered with this report include the generalizability of the current sample, which may be biased toward middle class, Caucasian, educated women from homogenous ethnic backgrounds. The cross-sectional design seriously limits the scope of the conclusions because the study design cannot address causation or the direction of correlational relationships. Volunteer bias may skew the sample toward more positive sexual attitudes and more sexual experience, and the use of retrospective self-report from questionnaires may reduce the accuracy of responses. A major limitation of this study was a lack of information regarding what type of genital pain women experienced (e.g., vulvar pain, deep pain, etc.). The study would have benefited from additional information, including the quality, location, and duration of pain, incorporation of pain rating scales for the estimation of pain intensity, and partner information to better understand the interpersonal effects of genital pain. Finally, the regression analyses employed can support statistical prediction but they are inappropriate for prospective prediction. Regression analyses can provide no information on causality or the direction of the relationship between sexual function, sexual satisfaction, and the predictor variables. Despite these concerns, the current study has provided insight into the impact of different levels of genital pain in young adult women. The sexual, behavioral, and attitudinal problems that emerged from this study suggest that the evaluation of levels of genital pain, rather than the presence or absence of pain, may be more useful in explaining the repercussions of painful intercourse.

**Acknowledgements** This publication was made possible by Grant Number 5 RO1 AT0022403 from the National Center for Complementary and Alternative Medicine to the second author. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of NCAAM or the NIH.

## References

- Barbanel, A. R. (1978). Diagnosis and treatment of coital discomfort. In J. LoPiccolo & L. LoPiccolo (Eds.), *Handbook of sex therapy* (pp. 241-259). New York: Plenum.
- Basson, R. (2001). Are the complexities of women's sexual function reflected in the new consensus definitions of dysfunction? *Journal of Sex & Marital Therapy*, 27(5), 112.
- Basson, R., Weijmar Shultz, W. C. M., Binik, Y. M., Brotto, L. A., Eschenbach, D. A., Laan, E., et al. (2004). Women's sexual desire and arousal disorders and sexual pain. In T. F. Lue, R. Basson, R. Rosen, F. Giuliano, S. Khoury, & F. Montorsi (Eds.), *Sexual medicine: Sexual dysfunctions in men and women* (pp. 851-974). Paris, France: Health Publications.
- Bergeron, S., Binik, Y. M., Khalife, S., & Pagidas, K. (1997). Vulvar vestibulitis syndrome: A critical review. *Clinical Journal of Pain*, 13(2), 42.
- Bergeron, S., Binik, Y. M., Khalife, S., Pagidas, K., Glazer, H. I., Meana, M., & Amsel, R. (2001). A randomized comparison of

- group cognitive-behavioral therapy, surface electromyography and biofeedback, and vestibulectomy in the treatment of dyspareunia resulting from vulvar vestibulitis. *Pain*, 91, 297-306.
- Binik, Y. M. (2005). Should dyspareunia be retained as a sexual dysfunction in DSM-V? A painful classification decision. *Archives of Sexual Behavior*, 34, 1-21.
- Brauer, M., Laan, E., & Kuile, M. (2006). Sexual arousal in women with superficial dyspareunia. *Archives of Sexual Behavior*, 35, 187-196.
- Bridges, S. K., Lease, S. H., & Ellison, C. R. (2004). Predicting sexual satisfaction in women: Implications for counselor education and training. *Journal of Counseling and Development*, 82, 158-166.
- Byers, E. S. (2005). Relationship satisfaction and sexual satisfaction: A longitudinal study of individuals in long-term relationships. *Journal of Sex Research*, 42, 113-118.
- Byers, E. S., & Demmons, S. (1999). Sexual satisfaction and sexual self-disclosure within dating relationships. *Journal of Sex Research*, 36, 180-189.
- Danielsson, I., Sjoberg, I., & Wikman, M. (2000). Vulvar vestibulitis: Medical, psychosexual and psychological aspects, a case-control study. *Acta Obstetrica et Gynecologica Scandinavica*, 79, 872-878.
- Davidson, J. K., & Moore, N. B. (1994). Masturbation and premarital sexual intercourse among college women: Making choices for sexual fulfillment. *Journal of Sex & Marital Therapy*, 19, 289-300.
- Davies, S., Katz, J., & Jackson, J. L. (1999). Sexual desire discrepancies: Effects on sexual and relationship satisfaction in heterosexual dating couples. *Archives of Sexual Behavior*, 28, 553-567.
- Davis, P., & Lay-Yee, R. (1999). Early sex and its consequences in New Zealand. *Journal of Sex Research*, 36, 135-144.
- Derogatis, L. R., & Melisaratos, N. (1979). The DSFI: A multidimensional measure of sexual functioning. *Journal of Sex & Marital Therapy*, 5, 244-281.
- Dove, N. L., & Wiederman, M. W. (2000). Cognitive distraction and women's sexual functioning. *Journal of Sex & Marital Therapy*, 26, 67-78.
- Dunne, M. P., & Najman, J. M. (2005). Is dyspareunia unrelated to early sexual abuse? *Archives of Sexual Behavior*, 34, 328-332.
- Friedman, M. (1995). Understanding vaginal vestibulum dyspareunia syndrome. *Cervix Lower Female Genital Tract*, 13, 35-40.
- Gates, E. A., & Galask, R. P. (2001). Psychological and sexual functioning in women with vulvar vestibulitis. *Psychosomatic Obstetrics and Gynecology*, 22, 11-28.
- Granot, M. (2005). Personality traits associated with perception of noxious stimuli in women with vulvar vestibulitis syndrome. *Journal of Pain*, 6, 168-173.
- Greiling, H., & Buss, D. (2000). Women's sexual strategies: The hidden dimension of extra-pair mating. *Personality and Individual Differences*, 28, 929-963.
- Haavio-Mannila, E., & Kontula, O. (1997). Correlates of increased sexual satisfaction. *Archives of Sexual Behavior*, 26, 99-119.
- Hill, C. A., & Preston, L. K. (1996). Individual differences in the experience of sexual motivation: Theory and measurement of dispositional sexual motives. *Journal of Sex Research*, 33, 27-45.
- Kaneko, K. (2001). Penetration disorder: Dyspareunia exists on an extension of vaginismus. *Journal of Sex & Marital Therapy*, 27, 153-155.
- Laumann, E. O., Paik, A., & Rosen, R. C. (1999). Sexual dysfunction in the United States: Prevalence and predictors. *Journal of the American Medical Association*, 281, 537-544.
- Lawrance, K., & Byers, E. S. (1995). Sexual satisfaction in long-term heterosexual relationships: The interpersonal exchange model of sexual satisfaction. *Personal Relationships*, 2, 267-285.
- Masheb, R. M., Lozano-Blanco, C., Kohorn, E. I., Minkin, M. J., & Kerns, R. D. (2004). Assessing sexual function and dyspareunia with the Female Sexual Function Index (FSFI) in women with vulvodynia. *Journal of Sex & Marital Therapy*, 30, 15-32.
- Meagher, M. W., Arnau, R. C., & Rhudy, J. L. (2001). Pain and emotion: Effects of affective picture modulation. *Psychosomatic Medicine*, 63, 79-90.
- Meana, M., Binik, Y. M., Khalife, S., & Cohen, D. R. (1997). Biopsychosocial profile of women with dyspareunia. *Obstetrics & Gynecology*, 90, 583-589.
- Meana, M., Binik, I., Khalife, S., & Cohen, D. (1998). Affect and marital adjustment in women's rating of dyspareunic pain. *Canadian Journal of Psychiatry*, 43, 381-385.
- Meana, M., Binik, Y. M., Khalife, S., & Cohen, D. (1999). Psychosocial correlates of pain attributions in women with dyspareunia. *Psychosomatics*, 40, 497-502.
- Mercer, C. H., Fenton, K. A., Johnson, A. M., Wellings, K., Maccowall, W., McManus, S., et al. (2003). Sexual function problems and help seeking behaviour in Britain: A national probability sample survey. *British Medical Journal*, 327, 426-427.
- Meston, C. M. (2003). Validation of the Female Sexual Function Index (FSFI) in women with female orgasmic disorder and in women with hypoactive sexual desire disorder. *Journal of Sex & Marital Therapy*, 29, 39-46.
- Meston, C., & Trapnell, P. (2005). Development and validation of a five-factor sexual satisfaction and distress scale for women: The Sexual Satisfaction Scale for Women (SSS-W). *Journal of Sexual Medicine*, 2, 66-81.
- Milhausen, R. R., & Herold, E. S. (1999). Does the sexual double standard still exist? Perceptions of university women. *Journal of Sex Research*, 36, 1-14.
- Nunns, D., & Mandal, D. (1997). Psychological and psychosexual aspects of vulvar vestibulitis. *Genitourinary Medicine*, 73, 541-544.
- Payne, K., Binik, Y. M., Amsel, R., & Khalife, S. (2004). Pain in women: An information processing approach to the study of vulvar vestibulitis syndrome. *Journal of Pain*, 5, 3-5.
- Pukall, C. F., Binik, Y. M., Khalife, S., Amsel, R., & Abbott, F. V. (2002). Vestibular tactile and pain thresholds in women with vulvar vestibulitis syndrome. *Pain*, 96, 163-175.
- Reissing, E. D., Binik, Y. M., Khalife, S., Cohen, D., & Amsel, R. (2003). Etiological correlates of vaginismus: Sexual and physical abuse, sexual knowledge, sexual self-schema, and relationship adjustment. *Journal of Sex & Marital Therapy*, 29, 47-59.
- Renaud, C., & Byers, E. S. (1997). Sexual and relationship satisfaction in mainland China. *Journal of Sex Research*, 34, 399-410.
- Rosen, R. C., Brown, C., Heiman, J., Leiblum, S., Meston, C. M., Shabsigh, R., et al. (2000). The Female Sexual Function Index (FSFI): A multidimensional self-report instrument for the assessment of female sexual function. *Journal of Sex & Marital Therapy*, 26, 191-208.
- Simpson, J. A., & Gangestad, S. W. (1991). Individual differences in sociosexuality: Evidence for convergent and discriminant validity. *Journal of Personality and Social Psychology*, 61, 870-883.
- Sprecher, S. (2002). Sexual satisfaction in premarital relationships: Associations with satisfaction, love, commitment, and stability. *Journal of Sex Research*, 39, 90-119.
- Sprecher, S., Barbee, A., & Schwartz, P. (1995). Was it good for you, too? Gender differences in first sexual intercourse experiences. *Journal of Sex Research*, 32, 1-15.
- Sprecher, S., & McKinney, K. (1993). *Sexuality*. Newbury Park, CA: Sage.
- van den Hout, M., & Barlow, D. (2000). Attention, arousal, and expectancies in anxiety and sexual disorders. *Journal of Affective Disorders*, 61, 241-256.

- Van Lankveld, J. J. D. M., Brewaeys, A. M. A., ter Kuile, M. M., & Weijnen, P. T. M. (1995). Difficulties in the differential diagnosis of vaginismus, dyspareunia, and mixed sexual pain disorder. *Journal of Psychosomatic Obstetrics & Gynaecology*, 16, 201-209.
- Van Lankveld, J. J. D. M., Weijnen, P. T. M., & ter Kuile, M. M. (1996). Psychological profiles of and sexual function in women with vulvar vestibulitis and their partners. *Obstetrics & Gynecology*, 88, 65-70.
- Vlaeyen, J. W., & Linton, S. J. (2000). Fear-avoidance and its consequences in chronic musculoskeletal pain: A state of the art. *Pain*, 85, 317-332.
- Waite, L. J., & Joyner, K. (2001). Emotional satisfaction and physical pleasure in sexual unions: Time horizon, sexual behavior, and sexual exclusivity. *Journal of Marriage and Family*, 63, 247-264.
- Wiegel, M., Meston, C. M., & Rosen, R. (2005). The Female Sexual Function Index (FSFI): Cross-validation and development of clinical cutoff scores. *Journal of Sex & Marital Therapy*, 31, 1-20.
- Wouda, J. C., Hartman, P. M., Bakker, R. M., Bakker, J. O., Van de Wiel, H. B. M., & Weijmar Schultz, W. C. M. (1998). Vaginal plethysmography in women with dyspareunia. *Journal of Sex Research*, 35, 141-147.